

MULTINETS, PARALLEL CONNECTIONS, AND MILNOR FIBRATIONS OF ARRANGEMENTS

Alex Suciu

Northeastern University

Joint work with Graham Denham (UWO)

Special Session on

Singular Spaces in Geometry, Topology, and Algebra

Spring Western Sectional Meeting

Boulder, Colorado

April 14, 2013

- 1 THE MILNOR FIBRATION OF AN ARRANGEMENT
- 2 CHARACTERISTIC VARIETIES
- 3 FINITE ABELIAN COVERS
- 4 ORBIFOLD FIBRATIONS
- 5 MULTINETS
- 6 MILNOR FIBERS
- 7 DELETION
- 8 POLARIZATION

THE MILNOR FIBRATION OF AN ARRANGEMENT

- Let \mathcal{A} be a (central) arrangement of hyperplanes in \mathbb{C}^{d+1} , with complement $M(\mathcal{A})$.
- Let $Q_m = \prod_{H \in \mathcal{A}} f_H^{m_H}$, where $H = \ker(f_H)$ and $m \in \mathbb{N}^{|\mathcal{A}|}$: a homogeneous polynomial of degree $N = \sum_{H \in \mathcal{A}} m_H$.
- The map $Q_m: \mathbb{C}^{d+1} \rightarrow \mathbb{C}$ restricts to a map $Q_m: M(\mathcal{A}) \rightarrow \mathbb{C}^*$.
- This is the projection of a smooth, locally trivial bundle, known as the (*global*) *Milnor fibration* of the multi-arrangement (\mathcal{A}, m) ,

$$F_m(\mathcal{A}) \longrightarrow M(\mathcal{A}) \xrightarrow{Q_m} \mathbb{C}^*.$$



- The typical fiber, $F_m(\mathcal{A}) = Q_m^{-1}(1)$, is called the *Milnor fiber* of the multi-arrangement.
- The (*geometric*) *monodromy* is the diffeomorphism

$$h: F_m(\mathcal{A}) \rightarrow F_m(\mathcal{A}), \quad z \mapsto e^{2\pi i/N} z.$$

- $F_m(\mathcal{A})$ is a Stein domain of complex dimension d ; thus, it has the homotopy type of a finite CW-complex of dimension d .
- If all $m_H = 1$, the polynomial $Q = Q_m$ is the usual defining polynomial, and $F(\mathcal{A}) = F_m(\mathcal{A})$ is the usual Milnor fiber of \mathcal{A} .

EXAMPLE

Let \mathcal{A} be a pencil of 3 lines through the origin of \mathbb{C}^2 . Then $F(\mathcal{A})$ is a thrice-punctured torus, and h is an automorphism of order 3:




- As conjectured by Papadima–S. and proved by Dimca–Papadima, Randell, Salvetti–Settepanella, etc, the complement $M(\mathcal{A})$ admits a minimal cell structure (Morse inequalities satisfied on the nose).
- In particular, $H_*(M(\mathcal{A}), \mathbb{Z})$ is torsion-free.
- Question (Dimca–Némethi, Randell): Is the same true for $F(\mathcal{A})$?
- In previous work (Cohen–Denham–S. 2003), we showed that the answer is no, but only for $F_m(\mathcal{A})$, for non-trivial m .
- Here, we settle the question for the standard Milnor fiber $F(\mathcal{A})$.

THEOREM

For every prime $p \geq 2$, there is a hyperplane arrangement whose Milnor fiber has non-trivial p -torsion in homology.

In particular, Milnor fibers of arrangements do not necessarily have a minimal cell structure.



Simplest example: the arrangement of **27** hyperplanes in \mathbb{C}^8 with

$$\begin{aligned}
 Q(\mathcal{A}) = & xy(x^2 - y^2)(x^2 - z^2)(y^2 - z^2)w_1w_2w_3w_4w_5(x^2 - w_1^2) \cdot \\
 & (x^2 - 2w_1^2)(x^2 - 3w_1^2)(x - 4w_1)((x - y)^2 - w_2^2)((x + y)^2 - w_3^2) \cdot \\
 & ((x - z)^2 - w_4^2)((x - z)^2 - 2w_4^2) \cdot ((x + z)^2 - w_5^2)((x + z)^2 - 2w_5^2).
 \end{aligned}$$

Then $H_6(F(\mathcal{A}), \mathbb{Z})$ has **2-torsion** (of rank **108**).

Basic idea of proof:

- $F_m(\mathcal{A})$ is the regular \mathbb{Z}_N -cover of $U = \mathbb{P}M(\mathcal{A})$ defined by the homomorphism $\delta_m: \pi_1(U) \rightarrow \mathbb{Z}_N$, taking each meridian x_H to m_H .
- If $\mathbb{k} = \overline{\mathbb{k}}$ and $\text{char}(\mathbb{k}) \nmid N$, then $H_q(F_m(\mathcal{A}), \mathbb{k}) \cong \bigoplus_{\rho} H_q(U, \mathbb{k}_{\rho})$, where $\rho: \pi_1(U) \rightarrow \mathbb{k}^*$ factors through δ_m .
- Thus, if there is such a character ρ for which $H_q(U, \mathbb{k}_{\rho}) \neq 0$, where $\rho = \text{char}(\mathbb{k}) \nmid N$, but there is no corresponding character in characteristic 0, then $H_q(F_m(\mathcal{A}), \mathbb{Z})$ will have ρ -torsion.
- Fix a prime p . Starting with an arrangement \mathcal{A} with a suitable multiset structure, we find a deletion $\mathcal{A}' = \mathcal{A} \setminus \{H\}$, and a choice of multiplicities m' on \mathcal{A}' such that $H_1(F_{m'}(\mathcal{A}'), \mathbb{Z})$ has p -torsion.
- Finally, we construct a "polarized" arrangement, $\mathcal{B} = \mathcal{A}' \parallel m'$, and show that $H_*(F(\mathcal{B}), \mathbb{Z})$ has p -torsion.

CHARACTERISTIC VARIETIES

- Let X be a connected, finite-type CW-complex. Then $G = \pi_1(X, x_0)$ is a finitely generated group, with $G_{\text{ab}} \cong H_1(X, \mathbb{Z})$.
- The character group $\widehat{G} = \widehat{G}_{\mathbb{k}} := \text{Hom}(G, \mathbb{k}^*) \cong H^1(X, \mathbb{k}^*)$ is an abelian algebraic group, with $\widehat{G} \cong \widehat{G}_{\text{ab}}$.
- Characteristic varieties:

$$\mathcal{V}_s^i(X, \mathbb{k}) := \{\rho \in \widehat{G} \mid \dim_{\mathbb{k}} H_i(X, \mathbb{k}_{\rho}) \geq s\}.$$

Here, \mathbb{k}_{ρ} is the local system defined by ρ , i.e., \mathbb{k} viewed as a $\mathbb{k}G$ -module, via $g \cdot x = \rho(g)x$, and $H_i(X, \mathbb{k}_{\rho}) = H_i(C_*(\tilde{X}, \mathbb{k}) \otimes_{\mathbb{k}G} \mathbb{k}_{\rho})$.

- Product formula: $\mathcal{V}_1^i(X_1 \times X_2, \mathbb{k}) = \bigcup_{p+q=i} \mathcal{V}_1^p(X_1, \mathbb{k}) \times \mathcal{V}_1^q(X_2, \mathbb{k})$.
- The sets $\mathcal{V}_s^1(X, \mathbb{k})$ depend only on $G = \pi_1(X)$ —in fact, only on G/G'' . Write them as $\mathcal{V}_s^1(G, \mathbb{k})$, and set $\mathcal{V}^1(G, \mathbb{k}) = \mathcal{V}_s^1(G, \mathbb{k})$.
- If $\varphi: G \rightarrow Q$ is an epimorphism, then the induced morphism $\hat{\varphi}_{\mathbb{k}}: \widehat{Q} \hookrightarrow \widehat{G}$ restricts to an embedding $\mathcal{V}_s^1(Q, \mathbb{k}) \hookrightarrow \mathcal{V}_s^1(G, \mathbb{k})$, $\forall s$.

EXAMPLE

We have $\widetilde{S^1} = \mathbb{R}$. Identify $\pi_1(S^1, *) = \mathbb{Z} = \langle t \rangle$ and $\mathbb{k}\mathbb{Z} = \mathbb{k}[t^{\pm 1}]$. Then:

$$C_*(\widetilde{S^1}, \mathbb{k}) : 0 \longrightarrow \mathbb{k}[t^{\pm 1}] \xrightarrow{t-1} \mathbb{k}[t^{\pm 1}] \longrightarrow 0.$$

For $\rho \in \text{Hom}(\mathbb{Z}, \mathbb{k}^*) = \mathbb{k}^*$, we get


$$C_*(\widetilde{S^1}, \mathbb{k}) \otimes_{\mathbb{k}\mathbb{Z}} \mathbb{k}_\rho : 0 \longrightarrow \mathbb{k} \xrightarrow{\rho-1} \mathbb{k} \longrightarrow 0,$$

which is exact, except for $\rho = 1$, when $H_0(S^1, \mathbb{k}) = H_1(S^1, \mathbb{k}) = \mathbb{k}$. Hence: $\mathcal{V}_1^0(S^1, \mathbb{k}) = \mathcal{V}_1^1(S^1, \mathbb{k}) = \{1\}$ and $\mathcal{V}_s^i(S^1, \mathbb{k}) = \emptyset$, otherwise.

EXAMPLE

Identify $\pi_1(\mathbb{C} \setminus \{n \text{ points}\}) = F_n$, and $\widehat{F}_n = (\mathbb{k}^*)^n$. Then:

$$\mathcal{V}_s^1(\mathbb{C} \setminus \{n \text{ points}\}, \mathbb{k}) = \begin{cases} (\mathbb{k}^*)^n & \text{if } s < n, \\ \{1\} & \text{if } s = n, \\ \emptyset & \text{if } s > n. \end{cases}$$



In general, though, the characteristic varieties of a space (or a group) will depend on the characteristic of the ground field.

EXAMPLE

- Let $\Gamma = F_n * \mathbb{Z}_p$, and identify $\widehat{\Gamma} = \widehat{F}_n \times \widehat{\mathbb{Z}}_p$.
- If $\text{char}(\mathbb{k}) \neq p$, then $\widehat{\mathbb{Z}}_p = \text{Hom}(\mathbb{Z}_p, \mathbb{k}^*)$ has order p , and

$$\mathcal{V}^1(\Gamma, \mathbb{k}) = \begin{cases} \widehat{\Gamma} & \text{if } n \geq 2, \\ (\widehat{\Gamma} \setminus \widehat{\Gamma}^\circ) \cup \{\mathbf{1}\} & \text{if } n = 1. \end{cases}$$

- If $\text{char}(\mathbb{k}) = p$, then $\widehat{\mathbb{Z}}_p = \{\mathbf{1}\}$, and $\mathcal{V}^1(\Gamma, \mathbb{k}) = \widehat{\Gamma}$, for all $n \geq 1$.

HOMOLOGY OF FINITE ABELIAN COVERS

- Let X be a connected, finite-type CW-complex, and $G = \pi_1(X)$.
- Let A be a finite abelian group.
- Every epimorphism $\chi: G \rightarrow A$ determines a regular, connected A -cover $X^\chi \rightarrow X$.
- Let \mathbb{k} be a field, $p = \text{char}(\mathbb{k})$. Assume $p = 0$ or $p \nmid |A|$. Then

$$H_q(X^\chi, \mathbb{k}) \cong H_q(X, \mathbb{k}[A]) \cong \bigoplus_{\rho \in \hat{A}} H_q(X, \mathbb{k}_\rho).$$

Hence,

THEOREM (LIBGOBER, SAKUMA, HIRONAKA)

$$\dim_{\mathbb{k}} H_q(X^\chi, \mathbb{k}) = \sum_{s \geq 1} |\mathcal{V}_s^q(X, \mathbb{k}) \cap \text{im}(\hat{\chi}_{\mathbb{k}})|,$$

where $\hat{\chi}_{\mathbb{k}}: \hat{A}_{\mathbb{k}} \rightarrow \hat{G}_{\mathbb{k}}$ is the induced morphism between character groups.

- Now suppose A is a finite cyclic group. Choose a generator α of A .
- Let $h = h_\alpha: X^\chi \rightarrow X^\chi$ be the monodromy automorphism.
- Let $h_*: H_q(X^\chi, \mathbb{k}) \rightarrow H_q(X^\chi, \mathbb{k})$ be the induced homomorphism.
- Note: if $\rho: G \rightarrow \mathbb{k}^*$ is a character belonging to $\text{im}(\hat{\chi}_\mathbb{k})$, there is a unique character $\iota_\rho: A \rightarrow \mathbb{k}^*$ such that $\rho = \iota_\rho \circ \chi$.

THEOREM

Assume $\text{char}(\mathbb{k}) \nmid |A|$. Then, the characteristic polynomial of the algebraic monodromy, $\Delta_{\chi, q}^{\mathbb{k}}(t) = \det(t \cdot \text{id} - h_*)$, is given by

$$\Delta_{\chi, q}^{\mathbb{k}}(t) = \prod_{s \geq 1} \prod_{\rho \in \text{im}(\hat{\chi}) \cap \mathcal{V}_s^q(X, \mathbb{k})} (t - \iota_\rho(\alpha)).$$

THEOREM

Let $X^\chi \rightarrow X$ be a regular, finite cyclic cover, defined by an epimorphism $\chi: \pi_1(X) \twoheadrightarrow \mathbb{Z}_r$. Suppose that $\text{im}(\hat{\chi}_{\mathbb{C}}) \not\subseteq \mathcal{V}_1^q(X, \mathbb{C})$, but $\text{im}(\hat{\chi}_{\mathbb{k}}) \subseteq \mathcal{V}_1^q(X, \mathbb{k})$, for some field \mathbb{k} of characteristic p not dividing r . Then $H_q(X^\chi, \mathbb{Z})$ has non-zero p -torsion.

- In order to apply this theorem, one needs to know both the characteristic varieties over \mathbb{C} and over \mathbb{k} , and exploit the qualitative differences between the two.
- In a special situation, we only need to verify that a certain condition on the first characteristic variety over \mathbb{C} holds.

THEOREM

Suppose there is a character $\rho: \pi_1(X) \rightarrow \mathbb{C}^*$ which factors as $\pi_1(X) \twoheadrightarrow \mathbb{Z} * \mathbb{Z}_p \twoheadrightarrow \mathbb{Z} \rightarrow \mathbb{C}^*$, for some prime p , but $\rho \notin \mathcal{V}^1(X, \mathbb{C})$. Then, for all sufficiently large integers r not divisible by p , there is a regular, r -fold cyclic cover $Y \rightarrow X$ such that $H_1(Y, \mathbb{Z})$ has non-zero p -torsion.

ORBIFOLD FIBRATIONS

- Let $\Sigma_{g,r}$ be a Riemann surface of genus $g \geq 0$ with $r \geq 0$ points removed. Fix points q_1, \dots, q_s on the surface, and assign integer weights μ_1, \dots, μ_s with $\mu_j \geq 2$.
- The orbifold $\Sigma = (\Sigma_{g,r}, \mu)$ is *hyperbolic* if $\chi^{\text{orb}}(\Sigma) := 2 - 2g - r - \sum_{i=1}^s (1 - 1/\mu_i)$ is negative.
- A hyperbolic orbifold Σ is *small* if either $\Sigma = S^1 \times S^1$ and $s \geq 2$, or $\Sigma = \mathbb{C}^*$ and $s \geq 1$; otherwise, Σ is *large*.
- Let $\Gamma := \pi_1^{\text{orb}}(\Sigma_{g,r}, \mu)$, defined as

$$\Gamma = \left\langle \begin{array}{c} x_1, \dots, x_g, y_1, \dots, y_g \\ z_1, \dots, z_s \end{array} \mid \begin{array}{l} [x_1, y_1] \cdots [x_g, y_g] z_1 \cdots z_s = 1 \\ z_1^{\mu_1} = \cdots = z_s^{\mu_s} = 1 \end{array} \right\rangle (r = 0)$$

$$\Gamma = F_{2g+r-1} * \mathbb{Z}_{\mu_1} * \cdots * \mathbb{Z}_{\mu_s} \quad (r > 0)$$

Then:

$$\mathcal{V}^1(\Gamma) = \begin{cases} \hat{\Gamma} & \text{if } \Sigma \text{ is a large hyperbolic orbifold,} \\ (\hat{\Gamma} \setminus \hat{\Gamma}^\circ) \cup \{1\} & \text{if } \Sigma \text{ is a small hyperbolic orbifold,} \\ \{1\} & \text{otherwise.} \end{cases}$$

- Let X be a smooth, quasi-projective variety, and $G = \pi_1(X)$.
- A surjective, holomorphic map $f: X \rightarrow (\Sigma_{g,r}, \mu)$ is called an *orbifold fibration* if the generic fiber is connected; the multiplicity of the fiber over each marked point q_i equals μ_i ; and f admits an extension $\bar{f}: \bar{X} \rightarrow \Sigma_g$ with the same properties.
- Such a map induces an epimorphism $f_{\#}: G \twoheadrightarrow \Gamma$, where $\Gamma = \pi_1^{\text{orb}}(\Sigma_{g,s}, \mu)$, and thus a monomorphism $\hat{f}_{\#}: \hat{\Gamma} \hookrightarrow \hat{G}$.

THEOREM (ARAPURA + ACM + BW)

$$\mathcal{V}^1(X) = \bigcup_{f \text{ large}} \text{im}(\hat{f}_{\#}) \cup \bigcup_{f \text{ small}} \left(\text{im}(\hat{f}_{\#}) \setminus \text{im}(\hat{f}_{\#})^{\circ} \right) \cup Z,$$

where Z is a finite set of torsion characters.

THEOREM

Suppose there is a small orbifold fibration $f: X \rightarrow (\Sigma, \mu)$ and a prime p dividing each μ_i . Then, for all $r \gg 0$ with $p \nmid r$, there is a regular, r -fold cyclic cover $Y \rightarrow X$ such that $H_1(Y, \mathbb{Z})$ has non-zero p -torsion.

MULTINETS

DEFINITION (FALK AND YUZVINSKY)

A (k, ℓ) -multinet ($k \geq 3, \ell \geq 1$) on an arrangement \mathcal{A} consists of:

- A partition $\mathcal{A} = (\mathcal{A}_1, \dots, \mathcal{A}_k)$.
- An assignment of multiplicities on the hyperplanes, $m: \mathcal{A} \rightarrow \mathbb{N}$.
- A subset $\mathcal{X} \subseteq L_2(\mathcal{A})$, called the base locus.

Moreover,

- $\sum_{H \in \mathcal{A}_i} m_H = \ell$, for all $i \in [k]$.
- For any hyperplanes H and H' in different classes, $H \cap H' \in \mathcal{X}$.
- For each $X \in \mathcal{X}$, the sum $n_X = \sum_{H \in \mathcal{A}_i: H \supset X} m_H$ is independent of i .
- For each $i \in [k]$, the space $(\bigcup_{H \in \mathcal{A}_i} H) \setminus \mathcal{X}$ is connected.

WLOG, we may assume $\gcd\{m_H \mid H \in \mathcal{A}\} = 1$. If all $m_H = 1$, the multinet is *reduced*. If, furthermore, every flat in \mathcal{X} is contained in precisely one hyperplane from each class, this is a *net*.




FIGURE : A $(3,2)$ -net on the A_3 arrangement




FIGURE : A $(3, 4)$ -multinet on the B_3 arrangement

- Let \mathcal{A} be a (central) arrangement of hyperplanes in \mathbb{C}^{d+1} .
- Let $M = \mathbb{C}^{d+1} \setminus \bigcup_{H \in \mathcal{A}} H$, and $U = \mathbb{P}M$. Then $M \cong U \times \mathbb{C}^*$.
- Identify $H_1(M, \mathbb{Z}) = \mathbb{Z}^n$, with basis the meridians $\{x_H \mid H \in \mathcal{A}\}$.
- Let $\text{Hom}(\pi_1(M), \mathbb{k}^*) = (\mathbb{k}^*)^n$ be the character torus.
- Then $\mathcal{V}^1(\mathcal{A}) := \mathcal{V}^1(M) \subset (\mathbb{k}^*)^n$ is isomorphic to $\mathcal{V}^1(U) \subseteq \{t \in (\mathbb{k}^*)^n \mid t_1 \cdots t_n = 1\} \cong (\mathbb{k}^*)^{n-1}$.

THEOREM (FALK-YUZ, PEREIRA-YUZ, YUZVINSKY)

Each positive-dimensional, non-local component of $\mathcal{V}^1(\mathcal{A})$ is of the form ρT , where ρ is a torsion character, $T = f^(H^1(\Sigma_{0,k}, \mathbb{C}^*))$, for some orbifold fibration $f: M(\mathcal{A}) \rightarrow (\Sigma_{0,k}, \mu)$, and either*

- $k = 2$, and f has at least one multiple fiber, or
- $k = 3$ or 4 , and f corresponds to a multinet with k classes on the multiarrangement (\mathcal{A}, m) , for some m .

MILNOR FIBERS

- Let (\mathcal{A}, m) be a multi-arrangement with $\gcd\{m_H \mid H \in \mathcal{A}\} = 1$. Set $N = \sum_{H \in \mathcal{A}} m_H$.
- The Milnor fiber $F_m(\mathcal{A})$ is the regular \mathbb{Z}_N -cover of $U(\mathcal{A})$ defined by the homomorphism $\delta_m: \pi_1(U(\mathcal{A})) \rightarrow \mathbb{Z}_N$, $x_H \mapsto m_H \bmod N$.
- If $\text{char}(\mathbb{k}) \nmid N$, then

$$\dim_{\mathbb{k}} H_q(F_m(\mathcal{A}), \mathbb{k}) = \sum_{s \geq 1} \left| \mathcal{V}_s^q(U(\mathcal{A}), \mathbb{k}) \cap \text{im}(\widehat{\delta}_m) \right|.$$

- The characteristic polynomial of the algebraic monodromy, $h_*: H_q(F_m(\mathcal{A}), \mathbb{k}) \rightarrow H_q(F_m(\mathcal{A}), \mathbb{k})$, is given by

$$\Delta_{h,q}^{\mathbb{k}}(t) = \prod_{s \geq 1} \prod_{\substack{\zeta \in \mathbb{k}^*: \zeta^N = 1, \\ \zeta^m \in \mathcal{V}_s^q(U(\mathcal{A}), \mathbb{k})}} (t - \zeta).$$

Not every regular, cyclic cover of a projective arrangement complement arises through the Milnor fiber construction. Nevertheless, the Milnor fibers dominate all other cyclic covers.

LEMMA

Let (\mathcal{A}, m) be a multiarrangement, and let $U^\chi \rightarrow U$ be a regular, r -fold cyclic cover of $U = \mathbb{P}M(\mathcal{A})$, given by $\chi: \pi_1(U) \rightarrow \mathbb{Z}_r$.

- $U^\chi \cong F_m(\mathcal{A})$ if and only if, for some integer k relatively prime to $N = \sum_{H \in \mathcal{A}} m_H$, the least positive integers μ_H satisfying $k \cdot \chi(x_H) = \mu_H \pmod N$ have the property that $\sum_{H \in \mathcal{A}} \mu_H = N$, in which case $\mu = m$.
- There exist infinitely many multiplicity vectors m for which the covering projection $F(\mathcal{A}, m) \rightarrow U$ factors through U^χ . Moreover, for any prime $p \nmid r$, we may choose m so that the degree of $Q(\mathcal{A}, m)$ is not divisible by p .

DELETION

A *pointed multinet* (\mathcal{M}, H) on an arrangement \mathcal{A} is a multinet structure $\mathcal{M} = ((\mathcal{A}_1, \dots, \mathcal{A}_k), m, \mathcal{X})$, together with a distinguished hyperplane $H \in \mathcal{A}$ for which $m_H > 1$ and $m_H \mid n_X$ for each $X \in \mathcal{X}$ such that $X \subset H$.


LEMMA

Suppose \mathcal{A} admits a pointed multinet, and set $\mathcal{A}' = \mathcal{A} \setminus \{H\}$. Then $U(\mathcal{A}')$ supports a small pencil, and $\mathcal{V}^1(\mathcal{A}')$ has a component which is a 1-dimensional subtorus, translated by a character of order m_H .

THEOREM

Suppose \mathcal{A} admits a pointed multinet, with distinguished hyperplane H and multiplicity m . Let p be a prime dividing m_H . There is then a choice of multiplicities m' on the deletion $\mathcal{A}' = \mathcal{A} \setminus \{H\}$ such that $H_1(F_{m'}(\mathcal{A}'), \mathbb{Z})$ has non-zero p -torsion.

POLARIZATION



- Let \mathcal{A}_1 and \mathcal{A}_2 be arrangements in V_1 and V_2 , defined by polynomials $Q_1 = \prod_{H \in \mathcal{A}_1} f_H$ and $Q_2 = \prod_{H \in \mathcal{A}_2} g_H$.
- Fix $H_1 \in \mathcal{A}_1$ and $H_2 \in \mathcal{A}_2$. The *parallel connection*, $\mathcal{A}_1 \parallel_{H_1, H_2} \mathcal{A}_2$, is the arrangement in $V_1 \times V_2$ defined by the polynomial

$$f_{H_1} \cdot \prod_{H \in \mathcal{A}_1 \setminus \{H_1\}} f_H \cdot \prod_{H \in \mathcal{A}_2 \setminus \{H_2\}} g_H \in \mathbb{k}[V_1^*] \otimes_{\mathbb{k}} \mathbb{k}[V_2^*] / (f_{H_1} - g_{H_2}).$$

- Falk and Proudfoot (2002): $\mathbb{P}(M_1) \times \mathbb{P}(M_2) \cong \mathbb{P}(M_1 \parallel_{H_1, H_2} M_2)$.

- If (\mathcal{A}_i, H_i) are pointed arrangements and $H \in \mathcal{A}_1$, we let

$$(\mathcal{A}_1, H_1) \circ_H (\mathcal{A}_2, H_2) = (\mathcal{A}_1_{H \parallel H_2} \mathcal{A}_2, H_1).$$

- Let $\mathcal{A} = (\{H_1, \dots, H_n\}, m)$ be a multi-arrangement. The polarization $\mathcal{A} \parallel m$ is the iterated parallel connection

$$\mathcal{A} \circ_{H_1} \mathcal{P}_{m_{H_1}} \circ_{H_2} \cdots \circ_{H_n} \mathcal{P}_{m_{H_n}},$$

where \mathcal{P}_k denotes a pencil of k lines in \mathbb{C}^2 .

- Note: $\text{rank}(\mathcal{A} \parallel m) = \text{rank } \mathcal{A} + |\{H \in \mathcal{A} : m_H \geq 2\}|$.
- Let $P_k = U(\mathcal{P}_k) \cong \mathbb{C} \setminus \{k - 1 \text{ points}\}$. Write $P(\mathcal{A}) = \prod_{H \in \mathcal{A}} P_{m_H}$. Then

$$U(\mathcal{A}) \times P(\mathcal{A}) \cong U(\mathcal{A} \parallel m).$$

Polarization construction works well with respect to Milnor fibrations:

$$\begin{array}{ccc}
 F_m(\mathcal{A}) & \longrightarrow & F(\mathcal{A} \parallel m) \\
 \downarrow & & \downarrow \\
 U(\mathcal{A}) & \longrightarrow & U(\mathcal{A} \parallel m),
 \end{array}$$




THEOREM

Suppose \mathcal{A} admits a pointed multinet, with distinguished hyperplane H and multiplicity m . Let p be a prime dividing m_H . There is then a choice of multiplicities m' on the deletion $\mathcal{A}' = \mathcal{A} \setminus \{H\}$ such that $H_q(F(\mathcal{B}), \mathbb{Z})$ has p -torsion, where $\mathcal{B} = \mathcal{A}' \parallel m'$ and $q = 1 + |\{K \in \mathcal{A}' : m'_K \geq 3\}|$.

EXAMPLE

- Let \mathcal{A} be reflection arrangement of type B_3 , with $Q(\mathcal{A}) = xyz(x-y)(x+y)(x-z)(x+z)(y-z)(y+z)$.
- Recall \mathcal{A} supports a $(3, 4)$ -multinet.
- Let $\mathcal{A}' = \mathcal{A} \setminus \{z = 0\}$ be the deleted B_3 arrangement.
- Pick $m' = (8, 1, 3, 3, 5, 5, 1, 1)$.
- Let $\mathcal{B} = \mathcal{A}' \parallel m'$, an arrangement of 27 hyperplanes in \mathbb{C}^8 .
- Then $H_6(F(\mathcal{B}), \mathbb{Z})$ has 2-torsion of rank 108.
- $\Delta_6^{\mathbb{C}}(t) = (t-1)^{11968}$, yet $\Delta_6^{\overline{\mathbb{F}_2}}(t) = (t-1)^{11968}(t^2 + t + 1)^{54}$.

REFERENCES

-  G. Denham, A. Suciu, *Multinets, parallel connections, and Milnor fibrations of arrangements*, arxiv:1209.3414.
-  A. Suciu, *Characteristic varieties and Betti numbers of free abelian covers*, arxiv:1111.5803, to appear in IMRN, 2013.
-  A. Suciu, *Hyperplane arrangements and Milnor fibrations*, arxiv:1301.4851.